

2021-06

# Complex photobiont diversity in the marine lichen *Lichina pygmaea*

Christmas, NAM

<http://hdl.handle.net/10026.1/18211>

---

10.1017/s002531542100062x

The Journal of the Marine Biological Association of the United Kingdom

Cambridge University Press

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

## Original Article

**Cite this article:** Christmas NAM, Allen R, Hollingsworth AL, Taylor JD, Cunliffe M (2021). Complex photobiont diversity in the marine lichen *Lichina pygmaea*. *Journal of the Marine Biological Association of the United Kingdom* **101**, 667–674. <https://doi.org/10.1017/S002531542100062X>

Received: 17 March 2021  
Revised: 24 August 2021  
Accepted: 24 August 2021  
First published online: 22 September 2021

### Key words:

Algae; cyanobacteria; host switching; intertidal zones; lichens; symbiosis

### Authors for correspondence:

Nathan Christmas, Michael Cunliffe,  
E-mail: [natchr@mba.ac.uk](mailto:natchr@mba.ac.uk), [micnli@mba.ac.uk](mailto:micnli@mba.ac.uk)

# Complex photobiont diversity in the marine lichen *Lichina pygmaea*

Nathan A. M. Christmas<sup>1</sup> , Ro Allen<sup>1</sup> , Anita L. Hollingsworth<sup>1,2</sup> ,  
Joe D. Taylor<sup>3</sup>  and Michael Cunliffe<sup>1,4</sup> 

<sup>1</sup>Marine Biological Association of the UK, The Laboratory, Citadel Hill, Plymouth, UK; <sup>2</sup>University of Southampton, European Way, Southampton, UK; <sup>3</sup>University of Bradford, Bradford, West Yorkshire, UK and <sup>4</sup>School of Biological and Marine Sciences, University of Plymouth, Plymouth, UK

## Abstract

Lichens are a well-known symbiosis between a host mycobiont and eukaryote algal or cyanobacterial photobiont partner(s). Recent studies have indicated that terrestrial lichens can also contain other cryptic photobionts that increase the lichens' ecological fitness in response to varying environmental conditions. Marine lichens live in distinct ecosystems compared with their terrestrial counterparts because of regular submersion in seawater and are much less studied. We performed bacteria 16S and eukaryote 18S rRNA gene metabarcoding surveys to assess total photobiont diversity within the marine lichen *Lichina pygmaea* (Lightf.) C. Agardh, which is widespread throughout the intertidal zone of Atlantic coastlines. We found that in addition to the established cyanobacterial photobiont *Rivularia*, *L. pygmaea* is also apparently host to a range of other marine and freshwater cyanobacteria, as well as marine eukaryote algae in the family Ulvophyceae (Chlorophyta). We propose that symbiosis with multiple freshwater and marine cyanobacteria and eukaryote photobionts may contribute to the ability of *L. pygmaea* to survive the harsh fluctuating environmental conditions of the intertidal zone.

## Introduction


Lichen symbiosis is undergoing renewed scrutiny, in part because of the application of molecular ecology techniques such as metabarcoding (Hawksworth & Grube, 2020; Smith *et al.*, 2020). Historic assumptions about the 'single fungus and single eukaryote alga or Cyanobacteria' nature of the symbiosis do not take into account tripartite lichens (i.e., those containing cyanobacterial and algal photobionts) (Rikkinen *et al.*, 2002; Henskens *et al.*, 2012), and have been further challenged by the more recent discovery of other fungi (Millanes *et al.*, 2016; Spribille *et al.*, 2016; Mark *et al.*, 2020), eukaryote algae (chlorobionts) and Cyanobacteria (cyanobionts) (Henskens *et al.*, 2012; Moya *et al.*, 2017) within lichens in addition to the primary myco- and photobionts (Smith *et al.*, 2020), alongside a complex diversity of heterotrophic bacteria (Grube *et al.*, 2009). This increased diversity of organisms co-existing within the lichen thallus has added support to an ongoing reinterpretation of lichens as even more complex ecological associations comprising numerous interacting components (Honegger, 1991; Hawksworth & Grube, 2020).

Amongst this complex lichen community, photobionts are fundamental members responsible for providing sugars (and in the case of some cyanobacterial photobionts, available nitrogen) essential for the survival of the lichen complex (holobiont). Extended photobiont diversity is likely widespread in terrestrial lichens (Dal Forno *et al.*, 2021) and has been hypothesized to be associated with increased ecological adaptive capability (Casano *et al.*, 2011; del Campo *et al.*, 2013; Muggia *et al.*, 2013; Ruprecht *et al.*, 2014). By recruiting a diverse range of photobionts, it has been proposed that lichens may benefit from enhanced ecological plasticity, facilitating the ability to survive a range of environments (Casano *et al.*, 2011; del Campo *et al.*, 2013; Muggia *et al.*, 2013). For example, the soil crust lichen *Psora decipiens* supports multiple lineages of *Trebouxia* and *Asterochloris* allowing it to dominate soil crusts from deserts to alpine sites (Ruprecht *et al.*, 2014). Additionally, photobiont 'switching' may allow for a further increase in fitness where multiple mycobiont lineages share locally adapted photobiont genotypes (Piercey-Normore & DePriest, 2001).

Changing environmental conditions are characteristic of the habitat occupied by marine lichens in the intertidal zone, with marine lichens being exposed to a uniquely dynamic range of pressures not experienced by terrestrial lichens including daily tidal cycles of seawater coverage. *Lichina pygmaea* (Lightf.) C. Agardh is a saxicolous fruticose lichen in the family Lichinaceae that inhabits the upper intertidal zone (Figure 1A). It can be found throughout the North Atlantic coastline, including the area surrounding Plymouth, UK (Naylor, 1930). *L. pygmaea* was historically thought to form a symbiosis with *Calothrix* (Cyanobacteria) (Whitton, 2012), which was updated through molecular characterization to *Rivularia* (Ortiz-Álvarez *et al.*, 2015). Marine lichens such as *L. pygmaea* remain relatively under-studied compared with terrestrial taxa, and the degree to which complex photobiont diversity occurs in marine lichens is yet to be determined. The potential for multiple cyanobionts within

© The Author(s), 2021. Published by Cambridge University Press on behalf of Marine Biological Association of the United Kingdom. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.





**Fig. 1.** (Ai–iii) *Lichina pygmaea* at Church Reef with close-up of thallus. Dotted line indicates approximate upper limit of the intertidal. Arrows point to *L. pygmaea* thalli. (B) Sampling locations of *L. pygmaea* near Plymouth (UK).

members of the Lichinaceae was proposed by Bubrick & Galun (1984), but the extent of this has not yet been investigated using molecular ecology techniques. Here, we perform a broad metabarcoding survey of the bacterial (16S rRNA gene) and eukaryote (18S rRNA gene) phototrophic taxa associated with the *L. pygmaea* thallus to establish the potential for complex photobiont diversity in a marine lichen.

## Methods

### Sampling

Sampling and processing were carried out according to the protocol described in Parrot *et al.* (2015). In summary, *L. pygmaea* thalli were collected from three locations close to Plymouth in the south-west of the UK (Figure 1B) in spring 2016: Rame Head (lat = 50.3228, long = −4.2194), West Wembury (lat = 50.3159, long = 4.0856), and Church Reef (lat = 50.3159, long = −4.0844). At each site, a single thallus was sampled using sterile forceps, transported to the laboratory at 4°C and processed within 1 h. Each of the three thalli were inspected using a dissecting microscope to remove any debris or invertebrates and separated into 1 g subsamples (N = 5). To separate the intra- and extrathalline communities, thalli were washed twice by placing in a universal tube with 20 mL sterile seawater and rotating for 10 min. Wash water was filtered through a 0.45 µm cellulose nitrate filter. Both the washed thalli (intrathalline community) and filter (extrathalline community) were homogenized using a Mini-Beadbeater machine (Biospec Products) at 42 rpm for 40 s.


### DNA extraction

DNA was extracted from the washed thalli (intrathalline) and the wash water filters (extrathalline) using the PowerSoil DNA Isolation Kit (MoBio, CA, USA) and stored at −20°C. The V4 region of the bacterial 16S rRNA gene was amplified using

primers 515F and 926R (Caporaso *et al.*, 2011) using Phusion® High-Fidelity DNA Polymerase (New England Biolabs) and PCR was performed under the following conditions: 94°C for 5 min, followed by 30 cycles of 94°C for 30 s, 53°C for 30 s and 72°C for 1 min, and a final elongation step at 5°C for 10 min. The V4 region of the eukaryotic 18S gene was targeted with the primer pair E572F and E1009R (Comeau *et al.*, 2011), and PCR was performed under the following conditions: 98°C for 30 s, followed by 30 cycles at 98°C for 10 s, 55°C for 30 s, 72°C for 30 s, with a final extension at 72°C for 5 min. Sequencing was performed on the Illumina MiSeq platform (Integrated Microbiome Resource, Dalhousie University, Canada).

### Bioinformatics

16S rRNA and 18S rRNA gene reads were processed through the DADA2 pipeline (Callahan *et al.*, 2016), which resolves amplicon sequence variants (ASVs) at single-nucleotide resolution (Callahan *et al.*, 2017). Reads were trimmed and truncated to remove primers and low-quality sequences respectively. Following error estimation, paired-end reads were merged, and chimeras were removed. Taxonomic classification of resolved ASVs was performed using the RDP naïve Bayesian classifier (Wang *et al.*, 2007) using SILVA version 132 (Quast *et al.*, 2013) for 16S rRNA gene ASVs and PR2 version 4.12.0 (Guillou *et al.*, 2013) for 18S rRNA gene ASVs. Read counts, taxonomic classifications, and sample metadata were compiled as a phyloseq object prior to downstream analysis (McMurdie & Holmes, 2013). *L. pygmaea* reads were removed from analysis. A median of 15,115 16S rRNA gene reads and 1551 processed 18S rRNA gene reads per sample were recovered (Raw sequencing data are deposited in the European Read Archive under ENA study PRJEB43698). Alpha- (Observed, Shannon Index, Pielou's Evenness) and beta-diversity (Bray–Curtis dissimilarity) analyses were carried out using phyloseq and differential abundances of



**Fig. 2.** (A) Relative abundance of bacterial and eukaryotic intrathalline and extrathalline communities. (B) Comparison of alpha-diversity between intrathalline (red) and extrathalline (blue) communities of Cyanobacteria and Ulvophyceae, indicating Observed diversity, Shannon Index and Pielou's Evenness. (C) Bray-Curtis NMDS plots showing beta-diversity of Cyanobacteria and Ulvophyceae intrathalline (red) and extrathalline (blue) communities. Shape indicates sampling site.

specific ASVs between the extra- and intrathalline samples were assessed using DESeq2 (Love *et al.*, 2014), based on non-normalized read count data and a significance threshold of  $\alpha = 0.05$ , with a value of +1 assigned to the matrix to allow for statistical comparison between samples where no reads were returned. This analysis allowed for a direct test of which ASVs are enriched within the thalli and therefore most likely to be acting as photobionts. All analyses were performed in the R environment (R Core Team, 2017).

### Phylogenetic trees

Identities of cyanobacterial (accession numbers MZ169867–MZ169909) and eukaryote algal (accession numbers MZ190002–MZ190020) ASVs identified by DESeq2 were confirmed by performing BLAST searches (megablast, excluding uncultured/environmental sample sequences,  $e$ -value = 0.05) against the NCBI nt database (9 November 2020) (see Supplementary material). Two 18S rRNA gene sequences (ASV\_433 and ASV\_251) had indeterminate identities and were removed from further analysis. To build phylogenetic trees, best BLAST hits for each ASV and an additional 2–5 close matches were obtained. Where possible, information about source material was recorded. *L. pygmaea* derived *Rivularia* sequences from Ortiz-Álvarez *et al.* (2015) were included in the cyanobacterial tree and photobionts of *Hydropuntaria maura* and *Wahlenbergiella striatula* from Thüs *et al.* (2011) in the algal tree. Sequences were aligned using MAFFT (using G-INS-i) and manually trimmed to remove gappy tails and poorly aligned regions. The 16S rRNA gene sequence ASV\_170 was poorly


aligned and excluded from further analysis. Phylogenies were reconstructed using the Cipres (Miller *et al.*, 2011) implementation of RAXML-HP (Stamatakis, 2014) using the GTR-CAT model and 1000 bootstrap replicates. Outgroups were specified for both Cyanobacteria (*Pseudanabaena*) and Ulvophyceae (*Trentepohlia*) trees. Resulting trees were first checked in FigTree (<https://github.com/rambaut/figtree>), before being annotated in R using ggtree (Yu *et al.*, 2017). All figures were manually edited for aesthetics in Inkscape (<https://inkscape.org/>).

### Results

Cyanobacteria dominated the bacterial community within all three thalli accounting for up to 49% of 16S rRNA gene reads (Figure 2A), while the green algae family Ulvophyceae (Chlorophyta) dominated the intrathalline eukaryotic community, accounting for up to 82% of the 18S rRNA gene reads (Figure 2A). Analyses of alpha-diversity indicated that despite increased abundance, Cyanobacteria exhibited significantly lower diversity (Shannon,  $P \leq 0.001$ ) and evenness (Pielou,  $P \leq 0.001$ ) inside the sampled thalli (Figure 2B). Similarly, Ulvophyceae showed a non-significant (Shannon,  $P = 0.0781$ ) reduction in diversity within the thalli and a significant reduction in evenness (Pielou,  $P \leq 0.005$ ) (Figure 2B). Analyses of beta-diversity revealed a clear distinction between the intrathalline and extrathalline communities of both Cyanobacteria (Adonis,  $P \leq 0.001$ ) and Ulvophyceae (Adonis,  $P \leq 0.001$ ) (Figure 2C).

A total of six dominant (average normalized read count >20) cyanobacterial ASVs were significantly enriched ( $P \leq 0.005$ ) within at least one location (Figure 3A). Three significantly





**Fig. 3.** Deseq2 plots indicating (A) Cyanobacteria and (B) Ulvophyceae ASVs enriched in the interior (red) and exterior (blue) of the *L. pygmaea* thallus. Prefix 'endo-' refers to intrathalline and 'exo-' refers to extrathalline. Circle size is the average of normalized read counts over all samples. Significant enrichments are labelled.

enriched ASVs, ASV\_6 (MZ169869), ASV\_18 (MZ169873) and ASV\_55 (MZ169883) accounting for up to 8.7, 4.6 and 1.5% of the intrathalline bacterial community respectively, and a fourth ASV\_73 (MZ169887), which was only found at Rame Head where it comprised up to 6.1% of the community, all belonged to *Rivularia*, the established *L. pygmaea* photobiont (Ortiz-Álvarez *et al.*, 2015). However, ASV\_2 (MZ169867) and ASV\_4 (MZ169868) (ASV\_4 was only found at West Wembury) which made up to 20.6% and 18.1% of the intrathalline bacterial community respectively were both related to *Pleurocapsa*, a genus not previously associated with *L. pygmaea*. Other cyanobacterial ASVs were derived from the genera *Acaryochloris*, *Phormidesmis* and *Synechocystis*, some of which showed increased abundance in the extrathalline community. Within the phylogenetic tree (100 sequences, total alignment length = 1465 bp), all *Rivularia* ASVs occurred within a clade containing mostly marine and saline water derived sequences, while the highly abundant *Pleurocapsa* ASV\_2 (MZ169867) fell within a freshwater dominated clade (Figure 4).


The two abundant ASVs (average normalized read count > 20) belonging to the Ulvophyceae were significantly enriched ( $P \leq 0.005$ ) in the intrathalline community (Figure 3B), ASV\_14 (MZ190003) (up to 16% intrathalline eukaryotes) and ASV\_7 (MZ190002) (up to 50% intrathalline eukaryotes). Phylogenetic analysis (72 sequences, total alignment length = 1163 bp) placed these within clades of marine derived *Paulbroadya* and *Blidingia* respectively (Figure 5).


## Discussion

Here we report a considerable diversity of cyanobacterial and algal lineages associated with the interior of the *L. pygmaea* thallus. Complex diversity of lichen cyanobionts such as this has not been widely reported, despite early indications from the Lichinaceae family (Bubrick & Galun, 1984). A previous molecular-based survey of *L. pygmaea* cyanobionts (Ortiz-Álvarez *et al.*, 2015), which used conventional PCR and Sanger sequencing only amplified the dominant cyanobiont *Rivularia*. The whole community metabarcoding approach used here (which detects all sequence variants even at low abundance) suggests that the diversity of cyanobacteria contained within the thallus of *L. pygmaea* may be more complex than previously considered (Ortiz-Álvarez *et al.*, 2015), and the existence of high diversity of photobionts within the lichen thallus such as shown

here has likely been overlooked in many lichens. Our findings mirror those of Onuț-Bränström *et al.* (2018) who discovered multiple *Trebouxia* genotypes in *Thamnolia* and *Cetraria* thalli following Ion Torrent sequencing, compared with a single genotype recovered by Sanger sequencing. More widespread use of community metabarcoding approaches to investigate photobiont diversity may therefore reveal a greater diversity of photobionts and hitherto unrecognized lichen partnerships.

While many terrestrial tripartite lichens are well recognized (Rikkinen *et al.*, 2002; Hensgens *et al.*, 2012), to the authors' knowledge the presence of Ulvophyceae within the *L. pygmaea* thallus is the first documented occurrence of a putative chlorobiont in this marine species. This potentially places *L. pygmaea* amongst several other cyanolichens shown to also contain chlorobionts (Hensgens *et al.*, 2012). In particular, *L. pygmaea* appears to associate with *Blidingia*, a blade-like multicellular alga that forms an unusual association with *Turgidoscolum ulvae* (Verrucariaceae) whereby the mycobiont colonizes algal filaments (Pérez-Ortega *et al.*, 2018). The exact nature of the relationship of *L. pygmaea* and *Blidingia* is yet to be established, but hints toward a unique interaction that has been previously overlooked. The other potential photobiont, *Paulbroadya* is known to form symbioses with several other freshwater and marine lichens in the family Verrucariaceae (Thüs *et al.*, 2011; Gasulla *et al.*, 2019) (Figure 5). The relationship here between the *L. pygmaea*-associated *Paulbroadya* and the same genus associated with the crustose marine lichen *H. maura* indicates the potential for sharing of photobionts between marine lichens. Photobiont 'switching' can occur between closely and distantly related lichens occupying the same habitat (Piercey-Normore & DePriest, 2001; Dal Forno *et al.*, 2021), offering a fitness advantage through the acquisition of locally adapted photobionts, and is likely commonplace (Yahr *et al.*, 2006, 2004; Magain *et al.*, 2017). The potential for photobiont sharing in *L. pygmaea* is supported by the repeated occurrence of the same Ulvophyceae ASVs between sample sites, and comparisons between photobiont communities in *L. pygmaea* and *H. maura* from the same site are now necessary to establish whether the same algal genotypes occur within both species of lichen. Interestingly, this is in contrast to several of the cyanobacterial lineages, including *Rivularia*, which exhibited variation between sites indicating either reduced dispersal and/or vertical transmission of cyanobionts leading to subsequent localized population structuring.





**Fig. 5.** 18S rRNA gene maximum likelihood phylogeny of Ulvophyceae associated with *Lichina pygmaea* ASVs significantly enriched in the intrathalline samples are indicated with \*. Prefix 'endo-' refers to intrathalline (red circles) and 'exo-' refers to extrathalline (blue circles). Circle size is relative abundance. Node labels indicate bootstrap support values for main clades. Dashed lines next to clade names indicate paraphyletic clades.

The phylogenetic placement of many of the potential *L. pygmaea* photobionts identified in this study within clades dominated by marine-derived lineages suggests that the lichen draws from a pool of marine photobionts, supporting the concept that photobiont selection is influenced by prevailing environmental conditions (Piercey-Normore & Deduke, 2011). Although the full ecological significance of the apparent complex photobiont diversity in *L. pygmaea* is yet to be determined, a selection of photobionts that enhance survival under different conditions could be a key factor allowing the lichen to occupy the marine environment. Other studies have proposed that the presence of multiple coexisting photobionts with different physiological properties may contribute to increased fitness of the lichen holobiont (Yahr *et al.*, 2006; Casano *et al.*, 2011). Since *L. pygmaea* is able to photosynthesize both in air and submerged in seawater (Raven *et al.*, 1990), the ability of *L. pygmaea* to host a variety of photobionts may provide it with an ecological advantage with multiple marine-derived photobionts promoting survival in the intertidal

zone by photosynthesizing when in seawater, while the presence of some freshwater-derived photobionts may allow for increased rates of photosynthesis at low tide and during periods of high freshwater input (e.g. rain events).

Further research into the relationship between *L. pygmaea* and associated photobionts is now necessary, including determining biogeographic factors in symbiont recruitment, establishing relatedness of photobionts between different marine lichen species and the extent of photobiont sharing, and photobiont localization within the thallus. Determining how much these photobionts contribute to carbon and, in the case of the cyanobacteria, nitrogen fixation will be important for understanding their contribution to biogeochemical processes, and relevant physiological experiments may be carried out using growth chambers and other experimental approaches (Henskens *et al.*, 2012). RNA studies could help to quantify photobiont activity in relation to environmental parameters including seasonal, tidal and diel cycles. This work could contribute to our understanding of the



evolution of *L. pygmaea* and adaptation of the Lichinaceae to marine environments.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S002531542100062X>.

**Acknowledgements.** The authors wish to thank two anonymous reviewers whose comments significantly improved the quality of the manuscript.

**Financial support.** NC, RA and MC were supported by the European Research Council (ERC) MYCO-CARB project (grant agreement number 772584).

## References

- Bubrick P and Galun M (1984) Cyanobiont diversity in the Lichinaceae and Heppiaceae. *The Lichenologist* **16**, 279–287.
- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA and Holmes SP (2016) DADA2: high-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**, 581–583.
- Callahan BJ, McMurdie PJ and Holmes SP (2017) Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME Journal* **11**, 2639.
- Caporaso JG, Lauber CL, Walters WA, Berg-lyons D, Lozupone CA, Turnbaugh PJ, Fierer N and Knight R (2011) Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences USA* **108**, 4516–4522.
- Casano LM, del Campo EM, García-Breijo FJ, Reig-Armíñana J, Gasulla F, del Hoyo A, Guéra A and Barreno E (2011) Two *Trebouxia* algae with different physiological performances are ever-present in lichen thalli of *Ramalina farinacea*. Coexistence vs competition? *Environmental Microbiology* **13**, 806–818.
- Comeau AM, Li WKW, Tremblay J-É, Carmack EC and Lovejoy C (2011) Arctic Ocean microbial community structure before and after the 2007 record Sea Ice Minimum. *PLoS ONE* **6**, e27492.
- Dal Forno M, Lawrey J, Sikaroodi M, Gillevet PM, Schuettelpelz E and Lücking R (2021) Extensive photobiont sharing in a rapidly radiating cyanolichen clade. *Molecular Ecology* **30**, 1755–1776.
- del Campo EM, Catalá S, Gimeno J, del Hoyo A, Martínez-Alberola F, Casano LM, Grube M and Barreno E (2013) The genetic structure of the cosmopolitan three-partner lichen *Ramalina farinacea* evidences the concerted diversification of symbionts. *FEMS Microbiology Ecology* **83**, 310–323.
- Gasulla F, Guéra A, de los Ríos A and Pérez-Ortega S (2019) Differential responses to salt concentrations of lichen photobiont strains isolated from lichens occurring in different littoral zones. *Plant and Fungal Systematics* **64**, 149–162.
- Grube M, Cardinale M, de Castro JV, Müller H and Berg G (2009) Species-specific structural and functional diversity of bacterial communities in lichen symbioses. *ISME Journal* **3**, 1105–1115.
- Guillou L, Bachar D, Audic S, Bass D, Berner C, Bittner L, Boute C, Burgaud G, de Vargas C, Decelle J, Del Campo J, Dolan JR, Dunthorn M, Edvardsen B, Holzmann M, Kooistra WHCF, Lara E, Le Bescot N, Logares R, Mahé F, Massana R, Montresor M, Morard R, Not F, Pawlowski J, Probert I, Sauvadet A-L, Siano R, Stoeck T, Vault D, Zimmermann P and Christen R (2013) The protist ribosomal reference database (PR2): a catalog of unicellular eukaryote small sub-unit rRNA sequences with curated taxonomy. *Nucleic Acids Research* **41**, D597–D604.
- Hawthornth DL and Grube M (2020) Lichens redefined as complex ecosystems. *New Phytologist* **227**, 1281–1283.
- Hensgens FL, Green TGA and Wilkins A (2012) Cyanolichens can have both cyanobacteria and green algae in a common layer as major contributors to photosynthesis. *Annals of Botany* **110**, 555–563.
- Honegger R (1991) Functional aspects of the lichen symbiosis. *Annual Review of Plant Physiology and Plant Molecular Biology* **42**, 553–578.
- Love MI, Huber W and Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550.
- Magain N, Miadlikowska J, Goffinet B, Sérusiaux E and Lutzoni F (2017) Macroevolution of specificity in Cyanolichens of the Genus *Peltigera* Section *Polydactylon* (Lecanoromycetes, Ascomycota). *Systematic Biology* **66**, 74–99.
- Mark K, Laanisto L, Bueno CG, Niinemets Ü, Keller C and Scheidegger C (2020) Contrasting co-occurrence patterns of photobiont and cystobasidiomycete yeast associated with common epiphytic lichen species. *New Phytologist* **227**, 1362–1375.
- McMurdie PJ and Holmes S (2013) Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* **8**, e61217.
- Millanes AM, Diederich P and Wedin M (2016) Cyphobasidium gen. nov., a new lichen-inhabiting lineage in the Cystobasidiomycetes (Pucciniomycotina, Basidiomycota, Fungi). *Fungal Biology* **120**, 1468–1477.
- Miller MA, Pfeiffer W and Schwartz T (2011) The CIPRES science gateway: a community resource for phylogenetic analyses. In *Proceedings of the 2011 TeraGrid Conference: Extreme Digital Discovery*, TG '11. New York, NY: Association for Computing Machinery, pp. 1–8. <https://doi.org/10.1145/2016741.2016785>.
- Moya P, Molins A, Martínez-Alberola F, Muggia L and Barreno E (2017) Unexpected associated microalgal diversity in the lichen *Ramalina farinacea* is uncovered by pyrosequencing analyses. *PLoS ONE* **12**, e0175091.
- Muggia L, Vancurova L, Škaloud P, Peksa O, Wedin M and Grube M (2013) The symbiotic playground of lichen thalli – a highly flexible photobiont association in rock-inhabiting lichens. *FEMS Microbiology Ecology* **85**, 313–323.
- Naylor GL (1930) Note on the distribution of *Lichina confinis* and *L. pygmaea* in the Plymouth District. *Journal of the Marine Biological Association of the United Kingdom* **16**, 909–918.
- Onuț-Brännström I, Benjamin M, Scofield DG, Heiðmarsson S, Andersson MGI, Lindström ES and Johannesson H (2018) Sharing of photobionts in sympatric populations of *Thamnia* and *Cetraria* lichens: evidence from high-throughput sequencing. *Scientific Reports* **8**, 4406.
- Ortiz-Álvarez R, de los Ríos A, Fernández-Mendoza F, Torralba-Burrial A and Pérez-Ortega S (2015) Ecological specialization of two photobiont-specific maritime cyanolichen species of the genus *Lichina*. *PLoS ONE* **10**, e0132718.
- Parrot D, Antony-Babu S, Intertaglia L, Grube M, Tomasi S and Suzuki MT (2015) Littoral lichens as a novel source of potentially bioactive Actinobacteria. *Scientific Reports* **5**, 15839.
- Pérez-Ortega S, Miller KA and Ríos ADL (2018) Challenging the lichen concept: *Turgidoscylum ulvae* (Verrucariaceae) represents an independent photobiont shift to a multicellular blade-like alga. *The Lichenologist* **50**, 341–356.
- Piercey-Normore MD and Deduke C (2011) Fungal farmers or algal escorts: lichen adaptation from the algal perspective. *Molecular Ecology* **20**, 3708–3710.
- Piercey-Normore MD and DePriest PT (2001) Algal switching among lichen symbioses. *American Journal of Botany* **88**, 1490–1498.
- Quast C, Priesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J and Glöckner FO (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* **41**, D590–D596.
- R Core Team (2017) *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Raven JA, Johnston AM, Handley LL and McINROY SG (1990) Transport and assimilation of inorganic carbon by *Lichina pygmaea* under emersed and submersed conditions. *New Phytologist* **114**, 407–417.
- Rikkinen J, Oksanen I and Lohtander K (2002) Lichen guilds share related cyanobacterial symbionts. *Science* **297**, 357.
- Ruprecht U, Brunauer G and Türk R (2014) High photobiont diversity in the common European soil crust lichen *Psora decipiens*. *Biodiversity and Conservation* **23**, 1771–1785.
- Smith HB, Dal Grande F, Muggia L, Keuler R, Divakar PK, Grewe F, Schmitt I, Lumbsch HT and Leavitt SD (2020) Metagenomic data reveal diverse fungal and algal communities associated with the lichen symbiosis. *Symbiosis* **82**, 133–147.
- Sprille V, Tuovinen V, Resl P, Vanderpool D, Wolinski H, Aime MC, Schneider K, Stabenheiner E, Toome-Heller M, Thor G, Mayrhofer H, Johannesson H and McCutcheon JP (2016) Basidiomycete yeasts in the cortex of ascomycete macrolichens. *Science* **353**, 488–492.
- Stamatakis A (2014) RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313.
- Thüs H, Muggia L, Pérez-Ortega S, Favero-Longo SE, Joneson S, O'Brien H, Nelsen MP, Duque-Thüs R, Grube M, Friedl T, Brodie J, Andrew CJ, Lücking R, Lutzoni F and Gueidan C (2011) Revisiting photobiont diversity in the lichen family Verrucariaceae (Ascomycota). *European Journal of Phycology* **46**, 399–415.



- Wang Q, Garrity GM, Tiedje JM and Cole JR** (2007) Naïve Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* **73**, 5261–5267.
- Whitton BA** (ed.) (2012) *Ecology of Cyanobacteria II*. Dordrecht: Springer. <https://doi.org/10.1007/978-94-007-3855-3>.
- Yahr R, Vilgalys R and Depriest PT** (2004) Strong fungal specificity and selectivity for algal symbionts in Florida scrub *Cladonia* lichens. *Molecular Ecology* **13**, 3367–3378.
- Yahr R, Vilgalys R and DePriest PT** (2006) Geographic variation in algal partners of *Cladonia subtenuis* (Cladoniaceae) highlights the dynamic nature of a lichen symbiosis. *New Phytologist* **171**, 847–860.
- Yu G, Smith DK, Zhu H, Guan Y and Lam TT-Y** (2017) ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution* **8**, 28–36.